WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

The state of the s			
(51) International Patent Classification 6:		(11) International Publication Number: WO 97/08345	
C12Q 1/68, 1/70, C12P 19/34, C07H 21/04, C07K 16/00	A1	(43) International Publication Date: 6 March 1997 (06.02.97)	
(21) International Application Number: PCT/US96/14250 (22) International Filing Date: 28 August 1996 (28.08.96)		CH, DE, DK, ES, FI, FR, GB, GR, JE, IT, LU, MC, NL.	
(30) Priority Data: 60/002,963 30 August 1995 (3Q.08.95)	τ	Published With international search report.	

MEDICAL COLLEGE [US/US]; Valhalla, NY 10595 (US).

(72) Inventors; and (75) Inventors/Applicants (for US only): DARZYNKIEWICZ Zbigniew [US/US]; 37 Meadow Lane, Chappagua, NY 10514 (US). LI, Xun [CN/US]; 1635 Morning few Drive, Yorktown Heigths, NY 10598 (US). TRAGANOS, Frank [US/US]; 301 66th Street, New York, NY 10021 (US).

(71) Applicant (for all designated States except US): NEW YORK

(74) Agents: MacWRIGHT, Robert, S. et al.; Skadden, Arps, Slate, Meagher & Flom, 919 Third Avenue, New York, NY 10022 (US).

AND THE METHODS FOR LABELING DNA ENDS WITH HALOGENATED NUCLEOTIDES AND DETECTING SAME WITH ANTIBODIES

All Abstract

The invention pertains to the field of DNA detection for basic research, medical diagnostic testing, and forensic testing. Methods are The invention pertains to the held of DNA detection for basic research, medical diagnostic testing, and forensic testing. Methods are provided for the end labeling of DNA strands. The pNA strands are first incubated with a halogenated deoxynucleotide triphosphate, such as brominated deoxynufline triphosphate (BrdUTP), and an ending the management of the didition of the halogenated deoxynucleotide to the 3'OH hends of the DNA strand, such as a termingly dooxynucleotide transferase (TdT). The resulting modified DNA strands are then incubated with a labeled antibody, such as a fluorescentate monoclonal antibody, that specifically binds to the halogenated deoxynucleotide. The label is then detected, e.g., by flow cytometry. The methods have utility in detecting apoptosis, in detecting DNA synthesis and/or repair, and as general methods for end labeling of DNA.